arXiv:1107.3569v1 [q-bio.CB] 18 Jul 2011

A dynamical system model for the onset of collective behavior in social amoebae

Y. Charles Li and Yipeng Yang

Abstract. Based upon the splendid experimental result of Gregor et al. [**3**], we present a dynamical system model on the motion of individual amoebae. The dynamics of our model qualitatively agrees well with Gregor et al.'s experimental result on the development of collective behavior in social amoebae.

1. Introduction

Complexity is the current novelty in Science. Chaos as a part of complexity has proved to be an important concept. Besides chaos, another concept that is very elusive in current complexity pursuits is emerging [**5**]. Among the phenomena that may be termed emerging, collective behavior of individual elements is the most intriguing. The potential importance of collective behavior was stressed long ago [**1**]. Recent more advanced experiments on microorganisms have revealed impressive collective behavior via dynamical quorum sensing [**2**]. We are especially impressed by the elegant experiment of Gregor et al. [**3**], which can manipulate and monitor an individual amoeba in the social amoebae *Dictyostelium discoideum.* An amoeba can synthesize and release into the extracellular space, a chemoattractant named cAMP (cyclic adenosine 3',5'-monophosphate). When it is stimulated with extracellular cAMP, an amoeba responds by synthesizing and releasing more cAMP [**7**]. The extracellular cAMP guides the amoebae aggregation to form fruiting bodies. The main discovery of Gregor et al. [**3**] is that collective behavior originates from the specific amoeba which is stimulated with extracellular cAMP. Here we shall introduce a dynamical system model to simulate the motion of the amoebae aggregation. Using mathematical equations to describe biological or chemical phenomena has a long history [**4**] [**8**] [**6**], and has established itself an important scientific area.

2. The model

Now we shall describe our model. Let N be the total number of amoebae ($N = 180$ in [**3**]). Denote by x_n the location of the n-th amoeba in the space.

1991 *Mathematics Subject Classification.* Primary 92; Secondary 37.

Key words and phrases. Collective behavior, social amoebae *Dictyostelium discoideum*, cAMP..

©2011 (copyright holder)

The space can be 3-dimensional, 2-dimensional, or 1-dimensional. The dynamical system modeling the motion of the amoebae is given by

$$\ddot{x}_n = -\sum_{j\in J} \nabla_{\dot{x}_n} G(|\dot{x}_n - \dot{x}_{n_j}|) - \sum_{k\in K} \nabla_{x_n} H(|x_n - x_{n_k}|) + \delta_{nn*} F(t), \tag{2.1}$$

where '$\cdot$' denotes time derivative d/dt, $|\ |$ denotes the Euclidean vector length, ∇ denotes the gradient, G is an attractive potential, H is an attractive-repulsive potential, J and K represent the neighbors, the G and H terms represent the forces of the extracellular cAMP, $F(t)$ is the stimulating force exercised by injecting extracellular cAMP at a specific amoeba, and

$$\delta_{nn*} = 1 \text{ for } n = n*, \ = 0 \text{ for } n \neq n*;$$

where $n*$ denotes the specific amoeba. We are interested in the following examples for G, H and F:

$$\begin{aligned} G(r) &= \frac{\alpha}{a} r^a, \\ H(r) &= \frac{\beta}{b}\left[\left(\frac{\sigma}{r}\right)^b - \frac{b}{c}\left(\frac{\sigma}{r}\right)^c\right], \\ F(t) &= \begin{pmatrix} \gamma \\ \gamma \end{pmatrix} \cos t; \end{aligned}$$

where $(\alpha, \beta, \gamma, a, b, c, \sigma)$ are positive constants. The model now takes the specific form

$$\begin{aligned} \ddot{x}_n &= -\alpha \sum_{j\in J} |\dot{x}_n - \dot{x}_{n_j}|^{a-2} (\dot{x}_n - \dot{x}_{n_j}) \\ &+ \beta \sum_{k\in K} \left[\left(\frac{\sigma}{|x_n - x_{n_k}|}\right)^b - \left(\frac{\sigma}{|x_n - x_{n_k}|}\right)^c\right] \frac{x_n - x_{n_k}}{|x_n - x_{n_k}|^2} \\ &+ \delta_{nn*} \begin{pmatrix} \gamma \\ \gamma \end{pmatrix} \cos t. \end{aligned} \tag{2.2}$$

The setup for the numerical simulation of (2.2) is as follows: The space is the 2-dimensional plane,

$$\alpha = 50, \quad a = 3, \quad \beta = 1, \quad b = 2,$$
$$c = 1, \quad \sigma = 0.1, \quad \gamma = 2, \quad N = 100;$$

thus each x_n is a 2-vector. For any n,

$$\text{if } |\dot{x}_n - \dot{x}_{n_j}| \leq 4, \text{ then } j \in J;$$

similarly

$$\text{if } |x_n - x_{n_k}| \leq 4, \text{ then } k \in K.$$

The initial locations of the 100 amoebae are uniformly distributed in a circular chamber of radius 10. The initial velocities of the 100 amoebae are uniformly distributed in

$$[-0.001, 0.001] \times [-0.001, 0.001].$$

We use the Runge-Kutta scheme with time step length 0.01 to conduct the numerical simulation. The motion of the 100 amoebae modeled by (2.2) qualitatively agrees well with the experimental result of Gregor et al. [**3**], see Figures 1-6.



Figure 1. The locations of the amoebae at time $t = 0$.

References

[1] P. Anderson, More is different, *Science* **177, no.4047** (1972), 393-396.

[2] S. De Monte, et al., Dynamical quorum sensing: population density encoded in cellular dynamics, *Proc. Natl. Acad. Sci. USA* **104, no.47** (2007), 18377-18381.

[3] T. Gregor, et al., The onset of collective behavior in social amoebae, *Science* **328** (2010), 1021-1025.

[4] A. Hodgkin, A. Huxley, A quantitative description of membrane current and its application to conduction and excitation in nerve, *J. Physiol.* **117** (1952), 500-544.

[5] R. MacKay, Nonlinearity in complexity science, *Nonlinearity* **21** (2008), T273-T281.

[6] J. Murray, *Mathematical Biology:* I. An Introduction (2002), II. Spatial Models and Biomedical Applications (2003), Springer.

[7] K. Tomchik, P. Devreotes, Adenosine 3',5'-monophosphate waves in Dictyostelium Discoideum: a demonstration by isotope dilution fluorography, *Science* **212** (1981), 443-446.

[8] A. Turing, The chemical basis of morphogenesis, *Phil. Trans. Roy. Soc. London, series B* **237, no.641** (1952), 37-72.

Department of Mathematics, University of Missouri, Columbia, MO 65211, USA
E-mail address: `liyan@missouri.edu`

Department of Mathematics, University of Missouri, Columbia, MO 65211, USA
E-mail address: `yangyip@missouri.edu`



Figure 2. The locations of the amoebae at time $t = 100$.



Figure 3. The locations of the amoebae at time $t = 200$.



Figure 4. The locations of the amoebae at time $t = 300$.



Figure 5. The locations of the amoebae at time $t = 400$.



FIGURE 6. The locations of the amoebae at time $t = 500$.